

Bag dimension: H355*W265+bottom 60

KEEP OUT OF REACH OF CHILDREN

MANCOZEB GROUP M3 FUNGICIDE



KENCOZEB 800 WP

Active Ingredient	: Mancozeb	800g/Kg
Inert Ingredients	: Lignosulfonic acid calcium salt	92.2g/Kg
	: NNO	34.8 g/Kg
	: Hexamethylenetetramine.....	16.4 g/Kg
	: Sodium dodecyl sulfate.....	8.1 g/Kg
	: Phthalocyanine Blue.....	4.4 g/Kg
	: Diatomaceous earth.....	to make 1Kg

FPA Registration No : 780-053-6461
Batch No. :
Date of Expiry :
Date of Manufacture :

Net Weight:
1 KG



Registered with the Fertilizer And Pesticide Authority Pursuant to P.D 1144

Imported and Distributed by: Kenso Philippines Inc.
9F Filinvest One Building, Northgate Cyberzone, Alabang-Zapote Road Corner Northgate Avenue,
Filinvest City, Alabang, Muntinlupa 1781, Philippines



Directions For Use

Use **KENCOZEB 800 WP** at the rate of 3-6 tablespoon (30-60 grams) per 16 liters of water. Spray at 7-10 days interval. Sprays should be directed uniformly to all parts of the plants. Use shorter interval and higher dosage under severe disease conditions. **KENCOZEB 800 WP** is compatible with commonly used insecticides, miticides, fungicides and surfactants. Do not spray within 7 days before harvest. Do not enter the field within 2 days after spraying.

Fruits, Nuts, Grains and Ornamentals

Crops	Weeds Controlled	Dosage	Method of Applications
Banana	<i>Sigatoka, Black Leaf Streak</i>	3 - 6 tbsp per 16 Liters of water	Spray at 10-12 days interval
Grape	<i>Black Rot, Downy Mildew</i>		Spray before and after blooms. Continue spraying through the season. Stop spraying 60days before harvest
Mango	<i>Anthraxnose, Scab</i>		Spray after flowering. Repeat at 15 days interval.
Melon	<i>Anthraxnose, Downy Mildew, Scab</i>		Begin spray when runners start to form or at the first sign of disease . Do not apply within 5 days of harvest.
Papaya	<i>Fruit Rot</i>		Start spray during flowering. Direct spray to crown blossom area
Watermelon	<i>Anthraxnose, Downy Mildew, Scab</i>		Begin spray when runners start to form or at the first sign of disease . Do not apply within 5 days of harvest.
Peanut	<i>Cercospora Leaf spot</i>		Spray when disease appears. Repeat at 10-14 days interval. Stop spraying 14 days before harvest.
Corn (sweet, popcorn)	<i>Helminthosporium Leaf spot</i>		Start application when disease first appears. Repeat at 4-7 days interval. Do not apply within 7 days of harvest.
Cotton	<i>Damping-off</i>		Spray when disease appear
Rice	<i>Blast, Cercospora Leaf Spots, Helminthosporium Leaf Spot, Sheath Blight</i>		Treat seedlings twice starting 3-4 leaf stage. Repeat spray at flowering stages. Repeat application as needed.
Soybean	<i>Leaf Rust</i>		Begin spray during late bloom followed by 2 more spray according to rate.
Ornamentals	<i>Anthraxnose, Black Spot, Blight</i>		Apply twice weekly during blooming season

Vegetables

Crops	Weeds Controlled	Dosage	Method of Applications
Beans	<i>Anthraxnose, Downy Mildew, Rust</i>	3 - 6 tbsp per 16 Liters of water	Spray when disease appears. Do not apply within 5 days of harvest
Broccoli	<i>Alternaria Leaf Spot, Downy Mildew</i>		Protect and spray field when disease threatens appears. Remove residues by trimming and washing. Do not apply within 7 days of harvest
Cabbage/ Cauliflower	<i>Alternaria Leaf Spot, Downy Mildew</i>		Treat seedlings from emergence to transplanting at 5-6 days interval appears. In the field, spray at 7-10 days interval.
Celery	<i>Late Blight</i>		Apply at 3-5 days interval appears as per dosage. Do not apply at 14 days of harvest.
Cucumber	<i>Anthraxnose, Downy Mildew, Scab</i>		Spray when runner begins to form or when disease appears. Repeat at 7 days interval.
Lettuce	<i>Downy Mildew</i>		Spray when disease appears.
Eggplant	<i>Anthraxnose, Late Blight</i>		Spray when fruits appears. Repeat at 7-10 days interval.
Mugbean	<i>Anthraxnose, Downy Mildew, Rust</i>		Spray when disease appears.
Onion/Garlic	<i>Botrytis Leaf Spot, Downy Mildew, Purple Blotch</i>		Spray at 4-6 leaf stage or when disease appears. Repeat at 7-10 days interval
Peas	<i>Anthraxnose, Downy Mildew, Rust</i>		Spray when disease appears Do not apply within 5 days of harvest.
Pepper/Chili	<i>Anthraxnose</i>		Spray when disease appears
Potato	<i>Early and Late Blight</i>		Spray when disease appears Repeat at 7-10 days interval
Squash	<i>Anthraxnose, Downy Mildew, Scab</i>		Begin spray at the first sign of disease
Tomato	<i>Anthraxnose, Early and Late Blight, Septoria Leaf Spot</i>		Treat seedlings from emergence to transplanting. In the field, spray as per dosage rate at 7-10 days interval.

KAUKULANG PAG-IINGAT

Iimbak sa orihinal na lalagyan at isarang mabuti. Iwasang madikit sa balat o malanghap. Iwasang malagyan ng kemikal ang ilog, lawa, kanal o anumang pinanggagalingan ng tubig. Huwag gamitin ang produktong ityo kapag hindi nauunawaan ang wastong paggamit mg pag iingat.

PRECAUTIONS: Store in original container, tightly closed. Avoid skin contact and inhalation of dust or spray mist. Avoid contamination of any water supply with chemical or empty container.

SECONDARY HAZARD: Irritant to skin on multiple exposure

FIRST AID: If swallowed and patient is conscious, give 1 glass of water, induce vomiting by tickling back of throat with a finger or blunt instrument. Repeat procedure until vomit fluid is clear in appearance.
In case of skin contact, remove contaminated clothing and wash affected areas with plain soap and water.
If inhaled, remove patient from contaminated area and bring patient to fresh air. If breathing has stopped apply artificial respiration.
In case of eye contamination, flush eye with clean running water for at least 15 minutes.
Get medical attention.

NATIONAL POISON MANAGEMENT AND CONTROL CENTER Tel. No. (02) 8524-1078/ 587-2067

PROHIBITION / WARRANTY STATEMENT

It is a violation of FPA rules and regulations to use this product in a manner inconsistent with its label. No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liability resulting from improper handling or use or application.